

МІНІСТЕРСТВО ОСВІТИ І НАУКИ УКРАЇНИ
ОДЕСЬКИЙ НАЦІОНАЛЬНИЙ УНІВЕРСИТЕТ
імені І. І. МЕЧНИКОВА

MINISTRY OF EDUCATION AND SCIENCE
OF UKRAINE
ODESSA I. I. MECHNIKOV NATIONAL
UNIVERSITY

**Фізика
аеродисперсних
систем**

**Physics
of aerodisperse
systems**

ВИПУСК

55

VOLUME

Науковий збірник

Collection of papers

Засновано в 1969 р.

Founded in 1969

О д е с а
О Н У
2018

Приведено результати досліджень тепло- і масообміну в дисперсних системах при протіканні фазових і хімічних перетворень, механізмів утворення дисперсної фази, фізики горіння різноманітних речовин, критичні умови займання та погасання. Окремо розглядається газодинаміка процесів переносу дисперсної фази. Представлено результати досліджень електрофізики дисперсних систем і нелінійних процесів, які протікають в димовій плазмі.

Видання призначено для науковців, які вивчають процеси в аеродисперсних системах, зокрема аерозолях. Воно має бути корисним для студентів та аспірантів фізичних, фізико-технічних, енергетичних і хімічних факультетів, які спеціалізуються в області теплофізики дисперсних систем і фізики плазми, хімічної фізики та гідродинаміки.

Редакційна колегія:

д. ф.-м. н., професор
д. ф.-м. н., професор
д. ф.-м. н., професор
д. ф.-м. н., професор
Ph.D., професор
д. ф.-м. н., професор
д. ф.-м. н., професор
д. ф.-м. н., професор
Ph.D., професор
д. ф.-м. н., професор
д. ф.-м. н., професор
д. х. н., професор
к. ф.-м. н., доцент
к. ф.-м. н., доцент
к. ф.-м. н.

Калінчак В. В. (Україна) (головний редактор)
Алтоїз Б. А. (Україна)
Асланов С. К. (Україна)
Ассовський І. Г. (Росія)
Гриншпун С. А. (США)
Драган Г. С. (Україна)
Контуш С. М. (Україна)
Копит М. Х. (Україна)
Новак Войцех (Польща)
Полетасєв М. І. (Україна)
Шевчук В. Г. (Україна)
Еннан А. А.-А. (Україна)
Орловська С. Г. (Україна) (відп. секретар)
Черненко О. С. (Україна) (техн. секретар)
Дараків Д. С. (Україна) (техн. секретар)
Красотова Л. М. (Україна) (техн. секретар)

Друкується згідно з рішенням Вченої ради ОНУ
від 30 жовтня 2018 р.

Включено до Переліку наукових профільних видань України згідно з наказом Міністерства освіти і науки України № 1279 від 6 листопада 2014 року.

Адреса редакційної колегії:

65082, Одеса, вул. Пастера, 27,

Одеський національний університет імені І. І. Мечникова,
кафедра загальної фізики та фізики теплоенергетичних і хімічних процесів,
тел. 048 / 723-12-03, 723-62-27

Адреса в Інтернеті: <http://fas.onu.edu.ua>

E-mail: teplophys@onu.edu.ua, svetor25@gmail.com, chernalex@ukr.net

The journal publishes results of studies in the field of heat and mass transfer in disperse systems during phase changes and chemical reactions, mechanisms of dispersed phase formation. Combustion of solid, liquid and gaseous fuels is analyzed, critical conditions of ignition and extinction are considered. Gas-dynamic phenomena that determine a transport of dispersed phase are considered. Studies of electrophysics of disperse systems and non-linear processes in dusty plasma are presented. The journal is intended for scientists and engineers in field of disperse systems and, in particular, aerosols. Also it will be useful for students and PhD students of physical, physical-technical, energy and chemical faculties, specializing in thermal and plasma physics of dispersed systems, chemical physics and hydrodynamics.

Editorial Board:

Kalinchak V. V. (Ukraine)	(Editor-in-Chief)
Altoiz B. A. (Ukraine)	
Aslanov S. K. (Ukraine)	
Assovsky I. G. (Russia)	
Dragan G. S. (Ukraine)	
Ennan A. A.-A. (Ukraine)	
Grinshpun S. A. (USA)	
Kontush S. M. (Ukraine)	
Kopyt N. Kh. (Ukraine)	
Nowak Wojciech (Poland)	
Poletaev N. I. (Ukraine)	
Shevchuk V. G. (Ukraine)	
Orlovskaya S. G. (Ukraine)	(Assistant Editor)
Chernenko A. S. (Ukraine)	(technical Secretary)
Darakov D. S. (Ukraine)	(technical Secretary)
Krasotova L. N. (Ukraine)	(technical Secretary)

The journal is published on the decision of the Academic Council of
Odessa I. I. Mechnikov National University of 30 October 2018

The journal is included in the list of scientific specialized editions of Ukraine according to the Ministry of Education and Science of Ukraine order № 1279, November 6, 2014.

Address of the Editorial Board:

65082, Odessa, Paster str. 27,
Odessa I. I. Mechnikov National University,
Department of general physics and physics of thermoenergetical and chemical
processes
Phone 048 / 723-12-03, 723-62-27
Internet address: <http://fas.onu.edu.ua>
E-mail: teplophys@onu.edu.ua, svetor25@gmail.com, chernalex@ukr.net

ЗМІСТ

ТЕПЛОФІЗИКА ДИСПЕРСНИХ СИСТЕМ

- Шут М. І., Рокицький М. О., Рокицька Г. В., Шут А. М., Стасюк І. М.*
Теплофізичні властивості полімерних композиційних матеріалів на основі пентапласту і вуглецевих нанотрубок 8
- Щекатолина С. А., Жарюк В. Н.*
Обмен липидами между липопротеинами низкой и высокой плотности с учетом окисления липопротеинов низкой плотности 14
- Zhou Huiyu, Dragan G. S., Kutarov V. V., Galkin N. B., Filipova T. O.*
Universal description of the biofilms growth Dynamic in logistic model 30
- Popovskii A. Yu., Altoiz B. A., Butenko A. F.*
Structural characteristics of the epitropic liquid-crystalline layer of oleic acid 37

ТЕПЛОМАСООБМІН

- Черненко А. С., Корчагина М. Н., Калинин В. В.*
Зажигание и горение частиц древесного угля в азотно-кислородных смесях комнатной температуры. Часть II. Аналитическое обоснование 45
- Orlovskaya S. G., Liseanskaia M. V.*
The oxide structures formation on the surface of tungsten 54

ФІЗИКА ГОРІННЯ

- Стариков М. А., Хлебникова М. Е., Буланін Ф. К., Сидоров А. Е., Полетаев Н. И., Шевчук В. Г.*
Горение диспергированного жидкого и твердого топлива в присутствии электрических разрядов 59
- Калінчак В. В., Черненко О. С., Сікорський М. В., Бритавський Е. В., Стукалов С. А.*
Експериментальні дослідження безполум'яного горіння газових сумішей з домішками аміаку, водню та ацетону на платиновому дротіку 71
- Буланін Ф. К., Сидоров А. Е., Кіро С. А., Полетаев Н. И., Шевчук В. Г.*
Критические условия воспламенения газовзвеси энергоемких соединений 80

ГАЗОДИНАМІКА

*Buyadzhi V. V., Sofronkov A. N., Glushkov A. V., Khetselius O. Yu.,
Dubrovskaya Yu. V., Svinarenko A. A.*

New energy, angle momentum and entropy balance approach to modelling climate and macroturbulent atmospheric dynamics, heat and mass transfer at macroscale. I. General Formalism

95

*Sofronkov A.N., Khetselius O.Yu., Glushkov A.V.,
Buyadzhi V.V., Romanova A.V., Ignatenko A.V.*

New geophysical complex-field approach to modelling dynamics of heat-mass-transfer and ventilation in atmosphere of the industrial region

104

ЕЛЕКТРОФІЗИКА

Вишняков В. И., Киро С. А., Кабаненко Р.В.

Методика зондовой диагностики плазмы сварочного аэрозоля

112

Драган Г. С., Колесников К. В., Кутаров В. В.

Температурный профиль сферического объема плазмы с учетом рекомбинационных процессов

121

ФІЗИКА АЕРОЗОЛІВ

Вишняков В. И., Опря М. В., Эннан А. А.-А.

Влияние щелочной присадки в защитном газе на дисперсный состав сварочного аэрозоля

126

Шингарёв Г.Л., Контущ С.М., Скапцов А.С.

Малоинерционный способ управления полями температур и пересыщений в термодиффузионных камерах. Опыты по исследованию диффузиофореза

139

Киро С. А., Опря М. В., Эннан А. А.-А.

Влияние параметров режима электродуговой сварки в защитном газе на дисперсный и химический состав частиц сварочного аэрозоля

149

ПРИЛАДИ І МЕТОДИ ВІМІРЮВАННЯ

Контущ С.М., Ахмеров А.Ю., Мандель В.Е., Чурашов В.П.

Пневматический генератор струи монодисперсных капель

158

Іменний показчик

167

Правила для авторів

169

CONTENTS

THERMAL PHYSICS OF DISPERSE SYSTEMS

- Shut M.I., Rokytskyi M.O., Rokytska H.V., Shut A.M., Stasiuk I.M.*
Thermophysical properties of polymer composite materials based on penton and carbon nanotubes 8
- Tchekatulina S. A., Garyuk V. N.*
Lipid exchange between lipoproteins of low and high density taking into account the oxidation of low-density lipoproteins 14
- Zhou Huiyu¹, Dragan G.S.², Kutarov V.V.², Galkin N.B.², Filipova T.O.²*
Universal description of the biofilms growth dynamic in logistic model 30
- Popovskii A.Yu., Altoiz B.A., Butenko A.F.*
Structural characteristics of the epitropic liquid-crystalline layer of oleic acid 37

HEAT AND MASS TRANSFER

- Chernenko A.S., Korchagina M.N., Kalinchak V.V.*
Ignition and combustion of charcoal particles in nitrogen-oxygen mixtures at room temperature. Part II Analytic justification 45
- Orlovskaya S. G., Liseanskaia M. V.*
The oxide structures formation on the surface of tungsten 54

COMBUSTION PHYSICS

- Starikov M.A., Khlebnikova M.E., Bulanin Ph.K., Sidorov A.E., Poletaev N.I., Shevchuk V.G.*
Dispersed liquid and solid fuels combustion in an electrical discharge presence 59
- Kalinchak V.V., Chernenko A.S., Sikorsky M.V., Brytavsky E.V., Stukalov S.A.*
Experimental studies of the flameless combustion of gas mixtures with impurities of ammonia, hydrogen and acetone on a platinum wire 71
- Bulanin Ph., Sidorov A., Kiro S., Poletaev N., Shevchuk V.*
Critical conditions of energy intense compounds aerosuspensions ignition 80

GAS-DYNAMICS

Buyadzhi V.V., Sofronkov A.N., Glushkov A.V.,

Khetselius O.Yu., Dubrovskaya Yu.V., Svinarenko A.A.

New energy, angle momentum and entropy balance approach to modelling climate and macroturbulent atmospheric dynamics, heat and mass transfer at macroscale. I. General Formalism

95

Sofronkov A. N., Khetselius O. Yu., Glushkov A. V.,

Buyadzhi V. V., Romanova A. V., Ignatenko A. V.

New geophysical complex-field approach to modelling dynamics of heat-mass-transfer and ventilation in atmosphere of the industrial region

104

ELECTROPHYSICS

Vishnyakov V. I., Kiro S.A., Kabanenko R.V.

Technique of probe diagnostics in the welding fume plasma.

112

Dragan G.S., Kolesnikov K.V., Kutarov V.V.

Temperature profile of a spherical plasma volume, taking into account recombination processes

121

AEROSOLS PHYSICS

Vishnyakov V.I., Oprya M.V., Ennan A. A.-A.

Effect of alkaline additional agent in shielding gas on disperse composition of welding fume.

126

Shynharov G.L., Kontush S.M., Skaptsov A.S.

A low inertia way of controlling the temperature and supersaturations fields in the thermodiffusion chambers. Experiments for research of the diffusiophoresis.

139

Kiro S.A., Oprya M.V., Ennan A.A.

Influence of the gas metal arc welding regime parameters on the disperse and chemical composition of the welding fume particles

149

INSTRUMENTS AND MEASUREMENT METHODS

Kontush S.M., Akhmerov O.Yu., Mandel V.Yu., Churashov V.P.

Pneumatic jet generator of monodisperse drops

158

Name index..... 168

Rules for authors..... 171

ГАЗОДИНАМІКА

УДК 539.328

**Buyadzhii V. V., Sofronkov A. N., Glushkov A. V.,
Khetselius O. Yu., Dubrovskaya Yu. V., Svinarenko A. A.**

Odessa State Environmental University

E-mail: buyadzhivv@gmail.com, a_sofronkov@ukr.net

New energy, angle momentum and entropy balance approach to modelling climate and macroturbulent atmospheric dynamics, heat and mass transfer at macroscale. I. General Formalism

We present elements of a new non-stationary balance approach to modelling global mechanisms of climate and macro turbulent atmospheric low-frequency processes, including processes of heat-mass transfer at spatial and temporal macro scales, teleconnection effects etc. The main forming factor is a triplet of interacting solitons: "the planetary soliton of Hadley cells - the entire complex of atmospheric fronts - the Rossby soliton wave packet". The approach is based on the using balance relationships for entropy, energy and angular momentum, spectral theory of atmospheric macroturbulence, atmospheric moisture flow in further connection with the continuity of atmospheric circulation forms (teleconnection, genesis of fronts). Particular application is studying a spatial-temporal picture of the long-term atmospheric pollutants (including distribution of radionuclides after accidents at the nuclear power plant like Fukushima etc) in the atmosphere with accounting for the macro turbulent, circulation low-frequency processes.

Introduction. Understanding and quantitative description of global mechanisms in atmospheric low-frequency processes, teleconnection effects etc attracts a fundamental interest in a modern physics of climate and heat-and mass transfer in complex atmospheric system " [1-20]. It should be noted that at present time there are different, quite consistent approaches to modelling global atmospheric macroprocesses and respectively the methods of modelling temporal and spatial dispersion of different pollutants in atmosphere and other geospheres (look for example, [21-30]). One should mention such methods as MLDP0 (Modèle Lagrangien de Dispersion de Particules d'ordre 0), HYSPLIT (Hybrid Single-Particle Lagrangian Integrated Trajectory Model), NAME (Numerical Atmospheric-dispersion Modelling Environment), RATM (Regional Atmospheric Transport Model), FLEXPART (Lagrangian Particle Dispersion Model), model of the European Center for Medium-Range Weather Forecasts (ECMWF) and others [7,11,26,30]. Nevertheless, correct quantitative description of the global atmospheric processes, the heat-and mass transfer processes in an atmosphere, macromodelling dispersion of the pollutants in an atmosphere remains very actual and hitherto unsolved problem. Below we present the elements of a new approach, which is based on the entropy, energy, moment balance relationships for the global atmospheric low-frequency processes, theory of atmospheric macrotur-

bulence and circulation over the position of the front sections. The fundamental idea is that the main process-forming factor is a triplet of interacting solitons: "the planetary soliton of Hadley cells - the entire complex of atmospheric fronts - the Rossby soliton wave packet" [22-24].

Advanced balance approach to macroatmospheric processes and a triplet of the interacting Hadley cells – Rossby – atmospheric fronts solitons. Nowadays it has become preferable to use single-factor models for the diagnosis and prediction of the atmospheric macrocirculation. However, these models do not reveal a dynamics of the process, but are only a summary statement of the state. For example, the forecast of climate changes in these models can only be done by studying trends in heat balance maps. In our opinion, the "heat balance" should be replaced by the "entropy indicator" of the climate state. This index satisfies the equation of the entropy balance. Moreover, it contains a time derivative, which does not exist in the heat balance equation. The problem is only that the entropy balance is calculated in the low-frequency range of climate change. Such a low-frequency range can only be in planetary-scale processes that form global climate change. It is well known [1, 22, 23] that such processes include wave oscillations such as the planetary soliton - the Hadley cells. These cells attract a great interest. For example, the only so far discovered process with a two-year cycle is the "southern process with El Niño", which is deciphered on the Hadley cells. The change in the state of the Hadley cells on a global scale will be the basis for saturation of the entropy balance equation, the minimum of which should be expected at the characteristic middle of the stage of the climatic epoch. A change in the energy of a soliton can occur if a moment of its destruction appears at low-frequency interaction with another, long-existing wave process, also of the soliton type. As a result of this interaction, a third soliton must arise. The second and third solitons are known: the Rossby solitons (blocks emerging from the tropical belt), which basically determine the nature of the atmosphere's circulation and the corresponding weather complex, and the soliton of the atmospheric fronts[22]. On the fronts, up to 30% of the solar energy is realized in the processes of wet convection to an effect of the atmospheric moisture flow. The thermodynamic front solenoids are so energy-saturated that they are a natural barrier to energy dissipation into mesoprocesses, the horizontal scale of which is much smaller than the transverse dimension of the front. Moreover, the transfer of energy from the mesoprocesses of the surface layer to the side of larger waves by the mechanism of "negative viscosity" is not carried out through the front. It explains that the main climate-forming (or global macro circulation) factor is a triplet of the solitons: "the planetary soliton of Hadley cells - the whole complex of atmospheric fronts - the Rossby soliton wave packet". The entropy balance equation should calculate the balance in the complex of the soliton triplet without mutual energy exchange (the active phase of the climatic stage), whereas the increasing entropy corresponds to a mutual energy exchange between the elements of the soliton triplet. All three elements of the soliton triplet determine the dynamic change in the complex of weather, while the factors that form the heat balance (radiation and heat fluxes) are basically subject to the annual course of insolation and do not form low-frequency processes of a climate-like

dynamics. The triplet of these solitons can be parametrized by means of the following model.

We assume that each of these solitons can be analytically determined by solving the known Karteweg-de Vries equation:

$$V_t - 6V^2V_x + V_{xxx} = 0. \quad (1)$$

Here V is the velocity, the subscripts mean the derivative with respect to the corresponding coordinate. The asymptotic solution (1) for time $t \rightarrow \infty$ has the form:

$$V(x, t) = (3t)^{-1/3} r(0) \text{Ai}(Z) - (3t)^{-2/3} i \frac{r'(0)}{2} \text{Ai}'(Z) + O\left(\frac{1}{3t}\right), \quad (2)$$

where $Z = x(3t)^{-1/3}$; r – generalized parameter of the soliton wave radiation matrix; $\text{Ai}(Z)$ – Airy function. The solution of Eq.(2) can be represented in the low-frequency range in the vicinity of $x = 0$ as a series of Z . The soliton of Hadley cells, forming a circumpolar vortex, occupies an area comparable to the hemisphere of the Earth; the Rossby soliton occupies a space of an order of magnitude smaller, while the transverse dimension of the front soliton is only about 30 km. The Rossby soliton is manifested as a result of the low energy transfer of the nonlinear interaction of these two solitons. In view of the apparent incomparability of the geometric area dimensions of these solitons, it becomes necessary to replace the coordinate of the space (x) by the energy coordinate (E). In this coordinate, all three solitons are comparable. The E -coordinate is formed on the basis of spectral modes of N -soliton solutions. The N -soliton solution has the following structure for the Karteweg-de Vries equation:

$$\psi_N = \sum_{\mu=0,1} \exp\left(\sum_i \mu_i \left(\eta_i + i \frac{\pi}{2}\right) + \sum_{i \leq j} \mu_i \mu_j A_{ij}\right), \quad (3)$$

where $\eta_i = k_i Z - k_i^3 t$ – wave vector in coordinate (Z, t); μ_i, μ_j – weight coefficients of particular solutions of the kink and antikink type; A_{ij} – matrix of a soliton nonlinear wave radiation:

$$e^{A_{ij}} = \left(\frac{k_i - k_j}{k_i + k_j}\right)^2. \quad (4)$$

Operator $\sum_{i \leq j} \mu_i \mu_j A_{ij}$ in the solution (4) determines the energy transformation

from the spectrum of N -solutions with radiation, i.e. transfer of the wave energy between the solitons. The construction (2) - (4) is to be introduced into the equation of the entropy balance:

$$\rho \frac{\partial S}{\partial t} + \nabla \cdot \frac{\vec{J}_q}{T} = \frac{|P| \cdot \vec{Y}}{T} + \frac{\vec{J}_q \cdot \vec{X}_q}{T}, \quad (5)$$

where ρ – is the substantial density of the medium through which the entropy flux S is carried out; $\vec{J}_q$ – the entropy flux vector; T is the temperature of the weighted average convection condition in the frontal systems, consistent with the Hadley cells; $\vec{X}_q = \nabla T$ – entropy productivity; $\vec{Y}_q = \nabla x \vec{V} + 2\vec{\omega}$ – an absolute vortex; $|P|$ – antisymmetric deformation tensor in space. Note that the third term in equation (5) spe-

cifies the irreversible course of the process in time.

Equation (5) must be rewritten for the coordinate E , i.e. it must correspond to the Riemannian geometry of the N -dimensional phase space of the spectral modes of N -soliton solutions. The soliton of the Hadley cells and the front soliton describe the curvature of the Riemannian space in the form of axial vectors, whereas the Rossby soliton contains mainly polar vectors. As is known, the tensor products of a tensor on axial vectors by a tensor on polar vectors give, as a result, the tensor of polar vectors. This means that the Rossby soliton, selecting the wave energy of the planetary soliton of the circumpolar vortex and the soliton of the fronts, gradually destroys them, translating them into a soliton of polar vectors. But the soliton of the front has a constant energy supply from wet convection and after its regeneration on the Rossby waves restores the structure of the planetary soliton. The cycles of such regeneration serve as the basis for low-frequency oscillations in the macroprocesses (climate) forecast and substantially exceed the periods of the frequency sweep of the solution (1). The, the long-period oscillation cycles of the tensor density arise in the Riemannian space time, as the energy losses generated by the wave radiation from the planetary soliton to the Rossby waves are compensated by the frontal soliton at the expense of the energy of the hydrologic cycle carried out in it. The curvature of the Riemannian space must also ensure the irreversibility in time of the tensor $|P|$ in equation (5). The friction losses are removed by the same mechanism. The form of the metric tensor of the Riemannian space for this problem was published in [22]. The Rossby solitons exist relatively a little time to be important for climate prediction. However, the fact of the repeated existence of blocking processes during the year ensures the existence of a total time interval sufficient for a significant increase in the field of the tensor density. This will be the time period that gives the necessary frequency in the task of forecasting the macroatmospheric processes. Two main solitons give wave energy to either Rossby solitons or progressive waves of temperate latitudes. But waves of temperate latitudes have a tensor analogy in axial vectors and therefore can not affect either the planetary soliton or the fronts. The energy costs for wave radiation in their direction are compensated by the latent energy of the phase transitions in the solitons of the fronts. The energy of the fronts cannot cause an energy overflow of the planetary soliton or solitons and the Rossby waves, as a process is limited by the solar constant. This nonlinear oscillatory process is stable in time and can exist long enough in time and commensurate with the time interval of climatic stage. The initial conditions in the sense of the Cauchy problem do not exist, because the system of solitons must enter a steady oscillatory regime over a long time interval of prehistory.

Advanced angle momentum balance approach and spectral analogue of atmosphere dynamics equations in a low-frequency range. As some elements of our theory were in details presented earlier [23-25], we are limited only by the key aspects. An advanced non-stationary angular momentum balance equation of in the planetary dynamic movements of air masses is written in the following standard integral form [24]:

$$\begin{aligned} \frac{\partial}{\partial t} \int \rho M dV = & \int_{\phi_1}^{\phi_2} \int_0^H \int_0^{2\pi} \rho v M d\phi dz d\lambda + \int_0^H \int_{\phi_1}^{\phi_2} \int_0^{2\pi} (p_E^i - p_W^i) a \cos \phi dz d\phi d\lambda + \\ & + \int_{\phi_1}^{\phi_2} \int_0^{2\pi} \int_0^H \tau_0 a \cos \phi d\phi d\lambda 2\pi, \end{aligned} \quad (6)$$

where $M = \Omega a^2 \cos \phi + u a \cos \phi$ – angular momentum; Ω – the angular velocity of rotation of the Earth; a – radius of the Earth; ϕ – Latitude ($\phi_1 - \phi_2$ – separated latitudinal belt between the Arctic and polar fronts); λ – longitude; u, v – zonal and meridional components of the wind velocity; ρ – air density; V – the entire volume of the atmosphere in this latitude belt from sea level to the average height of the elevated troposphere waveguide - H (in notations [1] $H = \infty$); $p_E^i - p_W^i$ – the pressure difference between the eastern and western slopes of the i -th mountains; z – height above sea level; τ_0 – the shear stress on the surface. From the point of view of physics, the cycle of balance of angular momentum in the contact zones with the hydrosphere and lithosphere becomes a singularity. This singularity can be detected through the occurrence of zones of fronts and soliton-type front. Then the kernel of equation (1) can be defined in the density functional ensemble of complex velocity potential [23, 24]

$$f = \overline{v_\infty} z + \frac{1}{2\pi} \sum_{k=1}^n q_k \ln(z - a_k) + \frac{1}{2\pi} \sum_{k=1}^p \frac{M_k e^{\alpha_k i}}{z - c_k} - \frac{i}{2\pi} \sum_{k=1}^m \Gamma_k \ln(z - b_k) \quad (7)$$

and the complex velocity, respectively, will be

$$v = U + iV = \frac{df}{dz} = \overline{v_\infty} + \frac{1}{2\pi} \sum_{k=1}^n \frac{q_k}{z - a_k} - \frac{1}{2\pi} \sum_{k=1}^p \frac{M_k e^{\alpha_k i}}{(z - c_k)^2} - \frac{i}{2\pi} \sum_{k=1}^m \Gamma_k / (z - b_k), \quad (8)$$

where w – complex potential; v_∞ – complex velocity general circulation background (mainly zonal circulation); b_k – coordinates of vortex sources in the area of singularity; c_k – coordinates of the dipoles in the area of singularity; a_k – coordinates of the vortex points in areas of singularity; M_k – values of momenta of these dipoles; α_k – orientation of the axes of the dipoles; Γ_k, q_k – values of circulation in the vortex sources and vortex points, respectively.

Further we consider an advanced spectral analogue for equation of motion for dynamics of the atmosphere in the low frequency range [22-24]. Note that the macro-turbulent atmosphere equations are low-frequency ones in its basis and there is a lot of experience of their solving by spectral methods (see [27-29]). The method for calculating a turbulence spectra should be based on the standard tensor equations of turbulent tensions. One could write the system of equations for the Reynolds tensions, moments of connection of the velocity pulsations with entropy ones and the corresponding closure equations [23, 24]. The technique of using Reynolds tension tensors of the second rank is well known (for example, in the form of an analytical representation). The master equations (velocity x-, y-components U, V) with accounting the Coriolis force can be rewritten as (look [24]):

$$\begin{aligned} \frac{\partial V'^2}{\partial t} = & -\frac{i}{a} \left[\overline{V'^2 L_1(\bar{V})} + 2\bar{V} \overline{V' L_1(V')} + \overline{V'^2 L_1(V')} \right] - \\ & -\frac{i}{a} \left[L_2(\bar{V}) \overline{V' U'} + \bar{V} \overline{U' L_2(V')} + \bar{U} \overline{V' L_2(V')} + \overline{V' U' L_2(V')} \right] + \\ & + 4\omega i \cos \theta \overline{V'^2} + \frac{2i}{a} \overline{V' L_6(\Phi')}, \end{aligned} \quad (9a)$$

$$\begin{aligned} \frac{\partial U'^2}{\partial t} = & -\frac{i}{a} \left[\overline{V' U' L_3(\bar{U})} + \bar{V} \overline{U' L_3(U')} + \bar{U} \overline{V' L_3(U')} + \overline{V' U' L_3(U')} \right] - \\ & -\frac{i}{a} \left[\overline{U'^2 L_4(\bar{U})} + 2\bar{U} \overline{U' L_4(U')} + \bar{U} \overline{U'^2 L_4(U')} \right] - 4\omega i \cos \theta \overline{U'^2} + \frac{2i}{a} \overline{U' L_5(\Phi')}, \end{aligned} \quad (9b)$$

where, as usually:

$$L_j = \frac{\partial(\dots)}{\partial \theta} - (-1)^j \frac{i}{\sin \theta} \frac{\partial(\dots)}{\partial \lambda} + b_j \operatorname{ctg} \theta(\dots), \quad b_j=1, j=1,4; b_j=-1, j=2,3; b_j=0, j=5,6.$$

The simplified modelling supposes remaining only two operators, say, in the equation (9c): $\frac{\partial V' U'}{\partial t} = \frac{i}{a} \overline{V' L_6(\Phi')}$ expressing Φ' through ϕ complex potential of the velocity f , and the velocity components U, V – in terms of functions ψ of the same velocity potential. The important parameter of the turbulent processes is the kinetic energy of turbulent vortices $b^2 = \overline{u'_k u'_k}$, which can be found from the equation (with physical explanations of any term):

$$\frac{\partial b}{\partial t} + \frac{\partial u_k b^2}{\partial x_k} + \frac{\partial}{\partial x_k} \left(\overline{u'_k u'_i u'_j} + 2 \overline{u'_k p'} \right) = -2 \overline{u'_k u'_i} \frac{\partial u_i}{\partial x_k} - 2 \frac{g}{\theta_0} \overline{w' \theta'}. \quad (10)$$

Advec- tion	Turbulent diffusion	Effect of forces of ten- sion	Interaction of Reynolds tension and averaged mo- tion	Generation for ac- count for swimming forces
----------------	------------------------	----------------------------------	--	--

Here g is the magnitude of the acceleration vector due to the planet's gravity, θ_0 is the equilibrium potential temperature, θ', p' are departures from equilibrium values. The equations for the velocity's correlates are in details listed in [23,24]. Components of tensor of the turbulent tensions are (spectral modes of velocity field):

$$\hat{V}^2 = \sum_{k=1}^{\infty} \sum_{s=-k}^k V_{k,s} T_{1,s}^k \left(\sum_{q=1}^{\infty} \sum_{j=-q}^q V_{q,j} T_{1,j}^q \right) = \sum_{k=1}^{\infty} \sum_{s=-k}^k \sum_{q=1}^{\infty} \sum_{j=-q}^q V_{k,s} V_{q,j} \times \sum_{v=|k-q|}^{k+q} \sigma_{1,1,2}^{k,q,v} \sigma_{s,j,s+j}^{k,q,v} T_{2,s+j}^v = \overline{v'_1 v'_1} = b^2 \quad (11)$$

Equating the velocity components determined in the global circulation model and model (6), one could find spectral matching between the wave numbers that define the functional elements in the Fourier-Bessel series with the source element of a plane field theory. It provides principally new basis for the macrocirculation atmospheric processes description. Such an approach is a key block in the balance approach to modelling global mechanisms of macro turbulent atmospheric low-frequency processes etc. In the next paper we will present the concrete simplified scheme to calculating the balance of an energy and angular momentum, atmospheric macro-turbu-

lence with elements of a chaos, atmospheric moisture flow in connection with the continuity of atmospheric circulation forms.

References:

1. *Peixoto J.P., Oort A.H.* Physics of Climate – N.-Y.: AIP, 1992. – 520p.
2. *Matveev V.* Physics of Atmosphere.-Moscow:Nauka, 1982.
3. *Glansdorff P. and Prigogine I.* Thermodynamic theory of structure, stability and fluctuations- New York, John Wiley Sons. – 1971.
4. *Iribarne J. V., Cho H. R.,* Atmospheric Physics.-N.-Y., Reidel Pub.Co., 1980.
5. *Ghil M., Kimoto M., Neelin J.D.* Nonlinear dynamics and predictability in the atmospheric sciences//Rev.Geophys. – 1991. – Vol.29. – P.46–55.
6. *Wang C.* ENSO, climate variability, and the Walker and Hadley circulations// In: The Hadley Circulation: Present, Past, and Future.- Eds Diaz H.F. and Bradley R.S.-Berlin: Springer. – 2004. – P.131-164.
7. *Khetselius O.Yu.,* Forecasting evolutionary dynamics of chaotic systems using advanced non-linear prediction method// Dynamical Systems – Theory and Applications, Eds. J. Awrejcewicz, M. Kazmierczak, P. Olejnik, J. Mrozowski. – 2013. – Vol.1. –P.145-152.
8. *Kurbatskii A. F.* Computational modeling of the turbulent penetrative convection above the urban heat island in stably stratified environment // Journal of Applied Meteorology. 2001. – V. 40. – P.1748 -1761.
9. *Lu J., Araya S.P., Snyder W.H., Jr. Lawson R.E.* A Laboratory Study of the Urban Heat Island in a Calm and Stably Stratified Environment. Part I: Temperature Field; Part II: Velocity Field // Journal of Applied Meteorology. – 1997. – V.36. – P.1377-1402.
10. *Brandstater A., Swinney H.* Strange attractors in weakly turbulent Couette-Taylor flow//Phys. Rev. A. – 1987. – Vol. 35. – P. 2207–2220.
11. *Gubanova E., Glushkov A., Khetselius O., Bunyakova Yu., Buyadzhi V., Pavlenko E.,* New methods in analysis and project management of environmental activity: Electronic and radioactive waste.– Kharkiv: FOP.-2017. – 120P.
12. *Wittman, M. A. H., Charlton, A. J. and Polvani, L. M.* On the meridional structure of annular modes// J. Climate. – 2005. – Vol. 18. – P.2119-2122.
13. *Voss R. and Mikolajewicz U.* Long-term climate changes due to increased CO₂ concentration in the coupled atmosphere-ocean general circulation model ECHAM3/LSG// Climate Dynamics. – 2001. –Vol.17. – P.45–60
14. *Kang I., Lau K.* Principal modes of atmospheric circulation anomalies associated with global angular momentum fluctuations//J.Atm.Sci. – 1994. – Vol.51. – P.1194.
15. *Wang, C.* ENSO, climate variability, and the Walker and Hadley circulations. The Hadley Circulation: Present, Past, and Future (Springer). – 2004.
16. *Boer G.J., Sargent N.E.* Vertically integrated budgets of mass and energy for the globe // J. Atmos. Sci. – 1985. – Vol. 42. – P.1592-1613.
17. *Trenberth K.E., Stepaniak D.P., Caron J.M.* Interannual variations in the atmospheric heat budget // J. Geophys. Res. – 2002. – Vol.107. – P.4.1-4.15.

18. *Fyfe J., Boer G., Flato G.* Predictable winter climate in the North Atlantic sector during 1997–1999 ENSO cycle// *Geophys.Res.Lett.* – 1999. – Vol.26. – P.1601-1604
19. *Arakava A., Shubert W.H.* Interaction of cumulus cloud ensemble with the large-scale environment//*Journ. of Atm. Sci.* – 1974. – Vol.31. – P.674-701.
20. *Khokhlov V.N., Glushkov A.V., Tsenenko I.A.* Atmospheric teleconnection patterns and eddy kinetic energy content: wavelet analysis// *Nonlinear Proc. in Geophys.* – 2004. – Vol.11. – P.295-301.
21. *Rusov V.D., Glushkov A.V., Vaschenko V.N., Myhalus O.T., Bondartchuk Yu.A., Smolyar V.P., Linnik E.P., Mavrodiev S.C., Vachev B.I.* Galactic cosmic rays – clouds effect and bifurcation model of the Earth global climate. Part 1. Theory // *Journal of Atmospheric and Solar-Terrestrial Phys.* – 2010. – Vol.72. – P.498-508.
22. *Glushkov A.V., Efimov V.A., Kivganov A.F.* Modelling climate as a problem of interaction of the solitons triplete//*Meteorology, Climatology and Hydrology.* – 1999. – Issue 38. – P.3-8.
23. *Glushkov A.V.* Renorm-group and fractal approach to turbulence spectrum in planetary atmosphere system, “cosmic plasma – galactic cosmic rays”//*Ukrainian Hydrometeorology Journal* – 2013. – №12. – P.25-30.
24. *Glushkov A., Khetselius O., Ambrosov S., Bunyakova Yu., Mansarliysky V.* The use of microsystems technology "Geomath" to modeling the balance of the angular momentum of the Earth, atmospheric processes and parameters of radio waveguides: III. Nonstationary theory//*Sensor Electr. and Microsyst. Techn.* – 2013. – Vol.10, №1. – P.22-28.
25. *Glushkov A., Safranov T., Khetselius O., Ignatenko A., Buyadzhi V., Svinarenko A.* Analysis and forecast of the environmental radioactivity dynamics based on methods of chaos theory: General conceptions//*Environm. Problems.* – 2016. – Vol.1, №2. – P.115-120.
26. *Glushkov A.V., Buyadzhi V.V., Ponomarenko E.L.* Geometry of Chaos: Advanced approach to treating chaotic dynamics in some nature systems// *Proc. Intern. Geom. Center.* – 2014. – Vol.7(1). – P.24-30.
27. *Glushkov A.V.* *Relativistic Quantum Theory. Quantum mechanics of Atomic Systems.* –Odessa: Astroprint, 2008.
28. *Khetselius O.Yu.* *Hyperfine structure of atomic spectra.* – Odessa: Astroprint, 2008.
29. *Malinovskaya S V, Glushkov A V, Khetselius O Yu, Svinarenko A A, Mischenko E.V., Florko T.A.* Optimized perturbation theory scheme for calculating the interatomic potentials and hyperfine lines shift for heavy atoms in the buffer inert gas//*Int. Journ.of Quantum Chemistry.* – 2009. – Vol.109, Issue 14. – P.3325-3329.
30. *Korsakissok I., Mathieu A., Didier D.* Atmospheric dispersion and ground deposition induced by the Fukushima Nuclear Power Plant accident: A local-scale simulation and sensitivity study// *Atm.Env.* – 2013. – Vol.70. – P.267-279

**Будджи В.В., Софронков О.Н., Глушков О.В.,
Хецелиус О.Ю., Дубровська Ю.В., Свиначенко А.А.**

Новый балансовый по энтропии, энергии и кутовому моменту подход к моделированию климата та макротурбулентной динамики атмосферы, тепло-массо-переносу в макромасштабе. I. Загальний формалізм

АНОТАЦІЯ

Представлені елементи нового балансового підходу до моделювання глобальних механізмів кліматичних і макротурбулентних атмосферних низькочастотних процесів, у т.ч., процесів тепло-массо-переносу, ефектів телеконекції тощо. Основним процес-формулюючим фактором є взаємодія триплету солітонів: «планетарний солітон осередків Гадля – комплекс атмосферних фронтів – хвильової пакет солітонів Россбі». Підхід заснований на використанні балансових співвідношень по ентропії, енергії і кутового моменту, спектральній теорії атмосферної макротурбулентності і вологообороту у зв'язку із наступністю форм атмосферної циркуляції (телеконекція, генезис фронтів). Одним із застосувань підходу є моделювання просторово-часових полів дисперсії забруднювачів (напр., хімічних забруднювачів типу з'єднань з галогенами, радіонуклідів після аварій на АЕС типу Фукусіми і т.і.) в атмосфері з урахуванням макротурбулентних циркуляційних низькочастотних процесів.

**Будджи В.В., Софронков А.Н., Глушков А.В., Хецелиус О.Ю.,
Дубровская Ю.В., Свиначенко А.А.**

Новый балансовый по энтропии, энергии и угловому моменту подход к моделированию климата и макротурбулентной динамики атмосферы, тепло-массо-переноса в макромасштабе. I. Общий формализм

АННОТАЦИЯ

Представлены элементы нового балансового подхода к моделированию глобальных механизмов климатических и макротурбулентных атмосферных низкочастотных процессов, в т.ч., процессов тепло-массо-переноса, эффектов телеконекции и др. Основным процесс-формирующим фактором является взаимодействие триплет солитонов: «планетарный солитон ячеек Гадля – весь комплекс атмосферных фронтов – волновой пакет солитонов Россби». Подход основан на использовании балансовых соотношений для энтропии, энергии и углового момента, спектральной теории атмосферной макротурбулентности и влагооборота в связи с преобладанием форм атмосферной циркуляции (телеконекция, генезис фронтов). Одно из применений подхода включает моделирование пространственно-временных полей дисперсии загрязнителей (напр., загрязнителей типа соединений галогенов, радионуклидов после аварий на АЭС типа Фукусимы и т.д.) в атмосфере с учетом макротурбулентных циркуляционных низкочастотных процессов.

Наукове видання

**ФІЗИКА
АЕРОДИСПЕРСНИХ
СИСТЕМ**

Випуск 55

Науковий збірник

Російською, українською та англійською мовами

Головний редактор **В.В. Калінчак**

Підписано до друку 12.12.2018. Формат 60х84/16.
Умов.-друк. арк. 10.11. Тираж 100 прим.
Зам. № 1858.

Надруковано з готового оригінал-макету

Видавець і виготовлювач
Одеський національний університет
імені І.І. Мечникова

Свідоцтво суб'єкта видавничої справи ДК № 4215 від 22.11.2011 р.

Україна, 65082, м. Одеса, вул. Єлісаветинська, 12
Тел.: (048) 723 28 39. E-mail: druk@onu.edu.ua